

REPRESENTATIVE DATA SHEET

Sheep anti-human Factor XII (F.XII)

Whole IgG from antiserum

5 mg

Product #: SAFXII-IG

Lot #: XXXX

Expiry date: XXXX

Store at -10 to -20°C

1395 Sandhill Drive, Ancaster, Ontario, Canada L9C 4V5
905-304-9896 • 800-903-6020 • fax 905-304-9897

For Research Use Only.

Not for use in diagnostic procedures.

Description of Factor XII (F.XII)

Factor XII (F.XII, Hageman factor) is a 76 kDa, single chain glycoprotein produced in the liver. In plasma, F.XII circulates as a protease zymogen at a concentration of approximately 30 µg/ml (400 nM). Upon vascular injury F.XII binds to negatively charged extravascular surfaces such as cartilage and skin, which facilitate activation of the zymogen to the active serine protease. Cleavage of F.XII by kallikrein after residue Arg³⁵³ produces the enzyme αF.XIIa, consisting of a 28 kDa light chain containing the protease domain, and a 52 kDa heavy chain containing the anionic surface-binding domain. Substrates for surface bound F.XIIa include the zymogens prekallikrein (PK) and factor XI (F.XI) as well as the procofactor high-molecular weight kininogen (HK). The activation of these substrates results in positive feedback activation of F.XII. Further cleavage of αF.XIIa by kallikrein produces the 28 kDa fragment βF.XIIa (Hageman factor fragment). βF.XIIa has reduced procoagulant activity as it lacks the anionic surface-binding domain, but is capable of fluid-phase activation of PK, factor VII and complement C1. The activity of F.XIIa in plasma is regulated predominantly by C1-Inhibitor, with relatively minor contributions by α₂antiplasmin, α₂macroglobulin and antithrombin, even in the presence of therapeutic levels of heparin¹⁻³.

REFERENCES and REVIEWS

1. DeLa Cadena R, Watchfogle YT, Colman RW, in Hemostasis and Thrombosis, 3rd Edition, eds. RW Colman, J Hirsh, VJ Marder and EW Salzman, pp. 219-240, J.B. Lippincott Co., Philadelphia, 1994.
2. Tankersley DL, Alving BM, Finlayson JS; Preparation of βF.XIIa (Hageman Factor Fragment) from Human Plasma. Thrombosis Research 25 pp 307-317, 1982.
3. Pixley RA, Schapira M, Coleman RW; The regulation of Human Factor XIIa by Plasma Proteinase Inhibitors. JBC 260, pp 1723-1729, 1985.

Document OPI0086, rev 1

Product Specifications

Description:

Vial containing XXXX ml of whole IgG representing approximately 1 ml of antiserum. Total protein is 5 mg.

Format:

Whole IgG, clear liquid.

Host Animal:

Sheep

Immunogen:

Human Factor XII purified from plasma.

Concentration:

IgG concentration is XXXX mg/ml, determined by absorbance using an extinction coefficient (E^{1%₂₈₀}) of 13.4.

Buffer:

10 mM HEPES, pH 7.4, 150 mM NaCl, 50% (v/v) glycerol.

Storage:

Store between -10 and -20°C. Product will become viscous but will not freeze. Avoid storage in frost-free freezers. Keep vial tightly capped. Allow product to warm to room temperature and gently mix before use.

Specificity:

This antibody is specific for factor XII as demonstrated by immunoelectrophoresis and ELISA.

Applications:

Suitable as a source of antibodies to human factor XII.

Neutralizing activity:

Not determined

Species Cross Reactivity: (immunodiffusion vs. citrated plasma)

Human:	XXXX	Mouse:	XXXX	Rat:	XXXX
Rabbit:	XXXX	Pig:	XXXX	Dog:	XXXX

Related Products:

Cat #: **GAFXII-IG** Goat anti-human F.XII, IgG from antiserum
 Cat #: **GAFXII-AP** Goat anti-human F.XII, affinity purified IgG
 Cat #: **GAFXII-HRP** Goat anti-human F.XII, IgG-peroxidase conjugate
 Cat #: **SAFXII-AP** Sheep anti-human F.XII, affinity purified IgG
 Cat #: **FXII-EIA** Paired antibody set for ELISA of F.XII, 5 x 96 wells
 Cat #: **FXII-DP** Human plasma deficient in F.XII, immune depleted

Visit our site (www.affinitybiologicals.com) for details.

Limited Warranty: This product is warranted to perform in accordance with its labeling and literature. Affinity Biologicals Inc. disclaims any implied warranty of merchantability or fitness for any other purposes, and in no event will Affinity Biologicals Inc. be liable for any consequential damages arising out of aforesaid express warranty.

Manufactured in Canada by:
AFFINITY BIOLOGICALS INC.
 1395 Sandhill Drive
 Ancaster ON CANADA L9C 4V5
 Tel: (905) 304-9896
 (800) 903-6020
 Fax: (905) 304-9897
info@affinitybiologicals.com